

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of: He et al.

Attorney Docket No.: PF140C2

Application Serial No.: Unassigned

Art Unit: Unassigned

Filed: Herewith

Examiner: Unassigned

Title: Interleukin-1 Beta Converting Enzyme
Like Apoptosis Protease-3 and 4

PRELIMINARY AMENDMENT

Commissioner For Patents
Washington, D.C. 20231

Sir:

Prior to examination, applicants hereby request that the following amendment and remarks be entered into the above-identified application. Applicants submit concurrently herewith: (a) Submission of Substitute/Formal Drawings of Figures 1A-B, 2A-B, and 3A-C (7 sheets) with copies of drawings as originally filed with changes marked; (b) a Substitute Sequence Listing in both paper and computer-readable forms; and (c) in accordance with 37 C.F.R. § 1.121(c)(1)(ii), a Version of Amendments With Markings to Show Changes Made. Please amend the application as follows.

In the Specification:

On page 1, after the title, please insert

--This application is a continuation of, and claims priority under 35 U.S.C § 120 to, U.S. Patent Application No. 08/334,251, filed November 1, 1994, which is herein incorporated by reference in its entirety.--.

Please replace the paragraph beginning at page 4, line7, with the following written paragraph:

-- Figures 1A-B show the cDNA and corresponding deduced amino acid sequence of ICE-LAP-3. The polypeptide encoded by the amino acid sequence shown is the putative mature form of the polypeptide (minus the

102020 0925360

initial methionine residue), and the standard one-letter abbreviation for amino acids is used. --

Please replace the paragraph beginning at page 4, line 12, with the following written paragraph:

-- Figures 2A-B show the cDNA and corresponding deduced amino acid sequence of ICE-LAP-4. The polypeptide encoded by the amino acid sequence shown is the putative mature form of the polypeptide (minus the initial methionine residue). --

Please replace the paragraph beginning at page 4, line 16, with the following written paragraph:

-- Figures 3A-C show an amino acid sequence comparison between ICE-LAP-3, ICE-LAP-4, human ICE and the *C. elegans* cell death gene *ced-3*. Shaded areas represent amino acid matches between the different sequences. --

Please replace the paragraph beginning at page 4, line 20, with the following written paragraph:

-- Sequencing inaccuracies are a common problem when attempting to determine polynucleotide sequences. Accordingly, the sequences of Figures 1A-B and 2A-B are based on several sequencing runs and the sequencing accuracy is considered to be at least 97%. --

Please replace the paragraph beginning at page 4, line 25, with the following written paragraph:

-- In accordance with an aspect of the present invention, there are provided isolated nucleic acids (polynucleotides) which encode the mature polypeptides having the deduced amino acid sequence of Figures 1A-B and 2A-B or for the mature polypeptide encoded by the cDNA of the clones deposited as ATCC Deposit No. 75875 encoding ICE-LAP-3, and ATCC Deposit No. 75873 encoding ICE-LAP-4, which were deposited August 25, 1994. --

Please insert the following paragraph on page 4, after line 32:

--These deposits are biological deposits with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Virginia 20110-2209. Since the strains referred to are being maintained under the terms of the Budapest Treaty, they will be made available to a patent office signatory to the Budapest Treaty.--

Please replace the paragraph beginning at page 4, line 33, with the following written paragraph:

-- The polynucleotide encoding ICE-LAP-3 can be detected from human prostate, human endometrial tumor, human pancreatic tumor, human adrenal gland tumor and human tonsil. The full-length encoding ICE-LAP-3 was discovered in a cDNA library derived from human endometrial tumor. It is structurally related to the Interleukin-1 converting enzyme family. It contains an open reading frame encoding a protein of approximately 341 amino acid residues. The protein exhibits the highest degree of homology to *C. elegans* cell death gene *ced-3* which is a homolog of human interleukin-1 converting enzyme, with 68 % similarity and 43 % identity over the entire amino acid sequence. It should be pointed out that the pentapeptide QACRG is conserved and is located at amino acid position 184-188. --

Please replace the paragraph beginning at page 5, line 21, with the following written paragraph:

-- The polynucleotides of the present invention may be in the form of RNA or in the form of DNA, which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand. The coding sequence which encode the mature polypeptides may be identical to the coding sequence shown in Figures 1A-B and 2A-B or that of the deposited clones or may be a different coding sequence which coding sequence, as a result of the redundancy or degeneracy of the genetic code, encode the same mature polypeptides, and derivatives thereof, as the DNA of Figures 1A-B and 2A-B or the deposited cDNA. --

Please replace the paragraph beginning at page 5, line 33, with the following written paragraph:

-- The polynucleotides which encode for the mature polypeptides of Figures 1A-B and 2A-B or for the mature polypeptides encoded by the deposited cDNAs may include: only the coding sequence for the mature polypeptide; the coding sequence for the mature polypeptide and additional coding sequence; the coding sequence for the mature polypeptide (and optionally additional coding sequence) and non-coding sequence, such as introns or non-coding sequence 5' and/or 3' of the coding sequence for the mature polypeptide. --

Please replace the paragraph beginning at page 6, line 10, with the following written paragraph:

-- The present invention further relates to variants of the hereinabove described polynucleotides which encode for fragments, analogs and derivatives of the polypeptides having the deduced amino acid sequence of Figures 1A-B and 2A-B or the polypeptides encoded by the cDNA of the deposited clones. The variants of the polynucleotides may be naturally occurring allelic variants of the polynucleotides or non-naturally occurring variants of the polynucleotides. --

Please replace the paragraph beginning at page 6, line 18, with the following written paragraph:

-- Thus, the present invention includes polynucleotides encoding the same mature polypeptides as shown in Figures 1A-B and 2A-B or the same mature polypeptides encoded by the cDNA of the deposited clones as well as variants of such polynucleotides which variants encode for a fragment, derivative or analog of the polypeptides of Figures 1A-B and 2A-B or the polypeptides encoded by the cDNA of the deposited clones. Such nucleotide variants include deletion variants, substitution variants and addition or insertion variants. --

Please replace the paragraph beginning at page 6, line 27, with the following written paragraph:

-- As hereinabove indicated, the polynucleotides may have a coding sequence which is a naturally occurring allelic variant of the coding sequence shown in Figures 1A-B and 2A-B or of the coding sequence of the deposited clones. As known in the art, an allelic variant is an alternate form of a polynucleotide sequence which may have a substitution, deletion or addition of nucleotides, which does not substantially alter the function of the encoded polypeptides. The polynucleotides may also encode for a proprotein which is the mature protein plus additional 5' amino acid residues. --

Please replace the paragraph beginning at page 7, line 11, with the following written paragraph:

-- The present invention further relates to polynucleotides which hybridize to the hereinabove-described sequences if there is at least 70%, preferably at least 90%, and more preferably at least 95% identity between the sequences. The present invention particularly relates to polynucleotides which hybridize under stringent conditions to the hereinabove-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. The polynucleotides which hybridize to the hereinabove described polynucleotides in a preferred embodiment encode polypeptides which either retain substantially the same biological function or activity as the mature polypeptide encoded by the cDNAs of Figures 1A-B and 2A-B or the deposited cDNA(s). --

Please replace the paragraph beginning at page 8, line 4, with the following written paragraph:

-- The present invention further relates to ICE-LAP-3 and 4 polypeptides which have the deduced amino acid sequence of Figures 1A-B and 2A-B or which has the amino acid sequence encoded by the deposited cDNAs, as well as fragments, analogs and derivatives of such polypeptides. --

Please replace the paragraph beginning at page 8, line 9, with the following written paragraph:

-- The terms "fragment," "derivative" and "analog" when referring to the polypeptides of Figures 1A-B and 2A-B or that encoded by the deposited cDNA, means polypeptides which retain essentially the same biological function or activity as such polypeptides, and wherein derivatives include polypeptides with enhanced or reduced biological function. An analog includes a proprotein portion to produce active mature polypeptides. --

Please replace the paragraph beginning at page 8, line 20, with the following written paragraph:

-- The fragment, derivative or analog of the polypeptides of Figures 1A-B and 2A-C or that encoded by the deposited cDNAs may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature polypeptide for purification of the mature polypeptide. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein. --

Please replace the paragraph beginning at page 26, line 11, with the following written paragraph:

--The DNA sequence encoding ICE-LAP-3, ATCC # 75875, is initially amplified using PCR oligonucleotide primers corresponding to the 5' sequences of the processed ICE-LAP-3 protein (minus the signal peptide sequence) and the vector sequences 3' to the ICE-LAP-3 gene. Additional nucleotides corresponding to ICE-LAP-3 are added to the 5' and 3' sequences respectively. The 5' oligonucleotide primer has the sequence 5' GATCGGATCCATGCGTGCGGGGACACGGGTC 3' (SEQ ID NO:5) contains a Bam HI restriction enzyme site (underlined) followed by 18 nucleotides of ICE-LAP-3 coding sequence starting from the presumed terminal amino acid of the processed protein codon. The 3' sequence 5'

GTACTCTAGATCATTACCCCTGGTGGAGGAT 3' (SEQ ID NO:6) contains complementary sequences to an Xba I site (underlined) followed by 21 nucleotides of ICE-LAP-3. The restriction enzyme sites correspond to the restriction enzyme sites on the bacterial expression vector pQE-9 (Qiagen, Inc. 9259 Eton Avenue, Chatsworth, CA, 91311). pQE-9 encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter operator (P/O), a ribosome binding site (RBS), a 6-His tag and restriction enzyme sites. pQE-9 is then digested with Bam HI and Xba I. The amplified sequences are ligated into pQE-9 and are inserted in frame with the sequence encoding for the histidine tag and the RBS. The ligation mixture is then used to transform *E. coli* available from Qiagen under the trademark M15/rep 4 by the procedure described in Sambrook, J. et al., Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989). M15/rep4 contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis. Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG ("Isopropyl-B-D-thiogalactopyranoside") is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression. Cells are grown an extra 3 to 4 hours. Cells are then harvested by centrifugation. The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl. After clarification, solubilized ICE-LAP-3 is purified from this solution by chromatography on a Nickel-Chelate column under conditions that allow for tight binding by proteins containing the 6-His tag (Hochuli, E. et al., J. Chromatography 411:177-184 (1984)). ICE-LAP-3 (95% pure is eluted from the column in 6 molar guanidine HCl pH 5.0 and for the purpose of renaturation adjusted to 3 molar guanidine HCl, 100mM sodium phosphate, 10 mmolar glutathione (reduced) and 2 mmolar glutathione (oxidized). After

incubation in this solution for 12 hours the protein was dialyzed to 10 mmolar sodium phosphate.--

Please replace the paragraph beginning at page 28, line 1, with the following written paragraph:

-- The DNA sequence encoding ICE-LAP-4, ATCC # 75873, is initially amplified using PCR oligonucleotide primers corresponding to the 5' sequences of the processed ICE-LAP-4 protein (minus the signal peptide sequence) and the vector sequences 3' to the ICE-LAP-4 gene. Additional nucleotides corresponding to ICE-LAP-4 are added to the 5' and 3' sequences respectively. The 5' oligonucleotide primer has the sequence 5' GATCGATCCATGGAGAACTGAAAACCTCA 3' (SEQ ID NO:7) contains a Bam HI restriction enzyme site (underlined) followed by 18 nucleotides of ICE-LAP-3 coding sequence starting from the presumed terminal amino acid of the processed protein codon. The 3' sequence 5' GTACTCTAGATTAGTGATAAAAATAGAGTTC 3' (SEQ ID NO:8) contains complementary sequences to an Xba I site (underlined) followed by 21 nucleotides of ICE-LAP-4. The restriction enzyme sites correspond to the restriction enzyme sites on the bacterial expression vector pQE-9 (Qiagen, Inc. 9259 Eton Avenue, Chatsworth, CA, 91311). pQE-9 encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter operator (P/O), a ribosome binding site (RBS), a 6-His tag and restriction enzyme sites. pQE-9 is then digested with Bam HI and Xba I. The amplified sequences are ligated into pQE-9 and are inserted in frame with the sequence encoding for the histidine tag and the RBS. The ligation mixture is then used to transform *E. coli* available from Qiagen under the trademark M15/rep 4 by the procedure described in Sambrook, J. et al., Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989). M15/rep4 contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis. Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp

(100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ration of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG ("Isopropyl-B-D-thiogalactopyranoside") is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression. Cells are grown an extra 3 to 4 hours. Cells are then harvested by centrifugation. The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl. After clarification, solubilized ICE-LAP-4 is purified from this solution by chromatography on a Nickel-Chelate column under conditions that allow for tight binding by proteins containing the 6-His tag (Hochuli, E. et al., J. Chromatography 411:177-184 (1984)). ICE-LAP-4 (95% pure is eluted from the column in 6 molar guanidine HCL pH 5.0 and for the purpose of renaturation adjusted to 3 molar guanidine HCl, 100mM sodium phosphate, 10 mmolar gluathione (reduced) and 2 mmolar glutathione (oxidized). After incubation in this solution for 12 hours the protein was dialyzed to 10 mmolar sodium phosphate.--

Please replace the paragraph beginning at page 30, line 7, with the following written paragraph:

-- The DNA sequence encoding for ICE-LAP-3, ATCC # 75875, was constructed by PCR on the full-length ICE-LAP-3 using two primers: the 5' primer 5' GACTATGCGTGCGGGGACACGG 3' (SEQ ID NO:9) contains the ICE-LAP-3 translational initiation site ATG followed by 5 nucleotides of ICE-LAP-3 coding sequence starting from the initiation codon; the 3' sequence 5' AATCAAGCGTAGTCTGGGACGTCGTATGGGTATTACCCCTGGTGGAGGATTTG 3' (SEQ ID NO:10) contains translation stop codon, HA tag and the last 21 nucleotides of the ICE-LAP-3 coding sequence (not including the stop codon). Therefore, the PCR product contains the ICE-LAP-3 coding sequence followed by HA tag fused in frame, and a translation termination stop codon next to the HA tag. The PCR amplified DNA fragment was ligated with pcDNAI/Amp by blunt end ligation. The ligation mixture was transformed into E. coli strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037) the

transformed culture was plated on ampicillin media plates and resistant colonies were selected. Plasmid DNA was isolated from transformants and examined by restriction analysis for the presence of the correct fragment. For expression of the recombinant ICE-LAP-3, COS cells were transfected with the expression vector by DEAE-DEXTRAN method (J. Sambrook, E. Fritsch, T. Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989)). The expression of the ICE-LAP-3 HA protein was detected by radiolabelling and immunoprecipitation method (E. Harlow, D. Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, (1988)). Cells were labelled for 8 hours with ³⁵S-cysteine two days post transfection. Culture media were then collected and cells were lysed with detergent (RIPA buffer (150 mM NaCl, 1% NP-40, 0.1% SDS, 1% NP-40, 0.5% DOC, 50mM Tris, pH 7.5) (Wilson, I. Et al., Id. 37:767 (1984)). Both cell lysate and culture media were precipitated with a HA specific monoclonal antibody. Proteins precipitated were analyzed on 15% SDS-PAGE gels.--

Page 31, line 6, after "SDS-PAGE gels." delete the remainder of the page which is empty space.

Please replace the paragraph beginning at page 32, line 20, with the following written paragraph:

-- The DNA sequence encoding for ICE-LAP-4, ATCC # 75873, was constructed by PCR on the full-length ICE-LAP-4 using two primers: the 5' primer 5' ACCATGGAGAACACTGAAAAC 3' (SEQ ID NO:11) contains the ICE-LAP-4 translational initiation site ATG followed by 5 nucleotides of ICE-LAP-4 coding sequence starting from the initiation codon; the 3' sequence 5' AATCAAGCGTAGTCTGGGACGTCGGTATGGGTAGTGATAAAAATA GAGTTCTTT 3' (SEQ ID NO:12) contains translation stop codon, HA tag and the last 21 nucleotides of the ICE-LAP-4 coding sequence (not including the stop codon). Therefore, the PCR product contains the ICE-LAP-4 coding sequence followed by HA tag fused in frame, and a translation termination stop codon next to the HA tag. The PCR amplified DNA fragment was ligated with pcDNA1/Amp by blunt end ligation. The ligation mixture was

transformed into E. coli strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037) the transformed culture was plated on ampicillin media plates and resistant colonies were selected. Plasmid DNA was isolated from transformants and examined by restriction analysis for the presence of the correct fragment. For expression of the recombinant ICE-LAP-4, COS cells were transfected with the expression vector by DEAE-DEXTRAN method (J. Sambrook, E. Fritsch, T. Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989)). The expression of the ICE-LAP-4 HA protein was detected by radiolabelling and immunoprecipitation method (E. Harlow, D. Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, (1988)). Cells were labelled for 8 hours with ³⁵S-cysteine two days post transfection. Culture media were then collected and cells were lysed with detergent (RIPA buffer (150 mM NaCl, 1% NP-40, 0.1% SDS, 1% NP-40, 0.5% DOC, 50mM Tris, pH 7.5) (Wilson, I. Et al., Id. 37:767 (1984)). Both cell lysate and culture media were precipitated with a HA specific monoclonal antibody. Proteins precipitated were analyzed on 15% SDS-PAGE gels.--

Please replace the sequence listing at pages 35-41 of the specification with the Substitute Sequence Listing (pages 1-6) attached hereto.

In the Figures:

Please cancel Figure 1, without prejudice, and replace it with new Figures 1A-B, submitted herewith with the formal drawings.

Please cancel Figure 2, without prejudice, and replace it with new Figures 2A-B, submitted herewith with the formal drawings.

Please cancel Figure 3, without prejudice, and replace it with new Figures 3A-C, submitted herewith with the formal drawings.

REMARKS

A substitute sequence listing and substitute Figures are submitted herewith to correct the nucleotide and amino acid sequences originally presented in the ICE-LAP 3 sequences (shown in the sequence listing as SEQ ID NOs:1 and 2 and in Figures 1 and 3). These changes do not introduce new matter because the correct sequence was inherent to the originally sequenced plasmid clone in actual possession of the Applicants before the time the captioned application was filed. The analysis needed to determine the complete and correct sequences were well within the skill of the ordinary artisan, and such analysis would not have required undue experimentation.

These same corrections to the Figures were submitted and entered in related Application No: 08/462,969 (now U.S. Patent No. 6,087,150). Attached hereto as Exhibit A is a copy of the Amendment and accompanying Declaration of Craig Rosen under 37 C.F.R. § 1.132 with Exhibits A-E thereto requesting correction of the sequence listing and figures for the aforementioned application. In this amendment, Applicants demonstrate that the corrected sequences are inherently present in the deposited material and, because Applicants have demonstrated "chain of custody" for the material originally sequenced and the resequenced material, correction of the originally presented sequence herein is not new matter. Furthermore, support for the correction of such sequencing errors can be found in the specification at page 4, lines 20 – 24 which state:

"Sequencing inaccuracies are a common problem when attempting to determine polynucleotide sequences. Accordingly, the sequences of Figures 1 and 2 are based on several sequencing runs and the sequencing accuracy is considered to be at least 97%."

The corrections to SEQ ID NO:1 submitted herein do not constitute a change of greater than 3%, hence the corrected sequence is fully supported by the specification and is not new matter.

Further, the Sequence Listing filed herewith includes sequences (SEQ ID NOs:5-12) corresponding to the primer sequences set forth in the examples of the instant application. The specification is amended as set forth above to insert the appropriate SEQ ID NOs at the appropriate locations for such primer sequences.

The specification has been amended to add the cross reference to the related application.

The specification was amended to reflect the drawing designations provided for in the formal drawings.

The specification was further amended on page 5 to correct the location of the pentapeptide QACRG in SEQ ID NO:2.

The specification was further amended on page 5 to correct the spelling of "homolog".

Accordingly, no new matter has been added herein.

Statements Under 37 C.F.R. § 1.825(a) and (b)

In accordance with 37 C.F.R. § 1.825(a), the undersigned attorney for Applicants hereby states that sequence information contained in the Substitute Sequence Listing submitted herewith is completely supported by the specification as originally filed, as detailed above, and no new matter has been introduced.

In accordance with 37 C.F.R. § 1.825(b), the undersigned attorney for Applicants hereby states that the information in the paper copy of the Substitute Sequence Listing submitted herewith is identical to the information contained in the computer readable form of the Substitute Sequence Listing submitted herewith.

SUMMARY

Applicants respectfully request entry of the amendments and remarks contained in this paper. If the Examiner believes that a telephone conference would help facilitate entry of the herein amendments, she is encouraged to call Applicants' representative at the telephone number listed below.

Respectfully submitted,

Dated: July 2, 2001


Jonathan L. Klein (Reg. No. 41,119)
Attorney for Applicants

Human Genome Sciences, Inc.
9410 Key West Avenue
Rockville, MD 20850
Telephone: (301) 251-6015

JLK/LT/ba

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re: application of: He et al.

Serial No.: 08/462,969

Group Art Unit: 1652

Filed: June 5, 1995

Examiner: G. Bugaisky

For: Interleukin-1 β Converting Enzyme
Like Apoptosis Protease 3 and 4

Attorney Docket No.: PF140P1

AMENDMENT

Assistant Commissioner
For Patents
Washington, D.C. 20231
Sir:

This Amendment is supplemental to the amendment filed March 15, 1999. A Declaration of Craig Rosen Under 37 CFR §1.132 accompanies this Amendment. Please amend the application as follows.

In the Specification:

On page 5, line 20, please replace "hoholog" with --homolog--.

On page 5, line 24, please replace "259-263" with --184-188--.

Please replace the sequence listing with the sequence listing attached hereto.

In the Figures

Please replace Figures 1 and 3 with the substitute Figures 1 and 3 attached hereto.

In the Claims:

61. (Once amended) An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence shown as [residues 1 to 341 in]SEQ ID NO:2;
- (b) the amino acid sequence shown as [residues 2 to 341 in]SEQ ID NO:2, but lacking the N-terminal methionine residue;
- (c) at least 30 contiguous amino acid residues of SEQ ID NO:2;
- (d) the amino acid sequence of a fragment of the polypeptide shown as residues 1 to 341 in SEQ ID NO:2 wherein said polypeptide fragment has apoptosis inducing activity; and
- (e) an amino acid sequence encoded by a polynucleotide capable of hybridizing to the complement of a polynucleotide consisting of SEQ ID NO:1, when incubated together in 0.5 M NaPO₄, pH 7.4 and 7% SDS overnight at 65°C followed by washing twice at room temperature and twice at 60°C with 0.5X SSC, and 0.1% SDS, wherein a polypeptide consisting of such amino acid sequence retains apoptosis inducing activity.

64. (Once amended) The isolated polypeptide of claim 63 comprising the amino acid sequence shown as [residues 2 to 341 in]SEQ ID NO:2, but lacking the N-terminal methionine residue.

65. (Once amended) The isolated polypeptide of claim [64] 63 comprising the amino acid sequence shown as [residues 1 to 341 in]SEQ ID NO:2.

REMARKS

Claims 61(b), 64, and 65 were amended herein. Claims 1-60 were canceled previously. Accordingly, claims 61-77 are pending. Reconsideration of the objections and rejections made in the Official action mailed October 14, 1998 is respectfully requested in light of both the Amendment filed March 15, 1999 and the present Amendment.

I. Amendments

A substitute sequence listing and substitute Figures 1 and 3 are submitted herewith to correct nucleotide and amino acid sequence errors as originally presented in the ICE-LAP 3 sequences (shown in the sequence listing as SEQ ID NOS:1 and 2). These changes do not introduce new matter because the correct sequence was inherent to the originally sequenced plasmid clone in actual possession of the Applicants before the time the captioned application was filed.

The captioned application is directed to both polynucleotides and polypeptides which are chemical compounds. The nucleotide sequence is but an inherent property of the described polynucleotides. There is a line of chemical case law where applicants have been permitted to amend the specification to correct the formula of a chemical compound after an application's filing date, provided that it had been demonstrated that one of skill in the art would have appreciated that the applicant was in possession of the compound itself at the time of filing. The rationale is that the formula is an inherent property of the compound and thus amending the specification to correct the formula is not new matter. See *In re Nathan*, 140 U.S.P.Q. 601, 604 (C.C.P.A. 1964). Accord *Kennecott Corp. v. Kyocera Int'l, Inc.*, 5 U.S.P.Q.2d 1194, 1198 (Fed. Cir. 1987), *cert. denied*, 486 U.S. 1008 (1988) ("The disclosure in a subsequent patent application of an inherent property of a product does not deprive that product of the benefit of the earlier filing date.").

In the field of biotechnology, applicants often rely on a deposited clone, where the deposit was made prior to filing, to establish possession of nucleic acids or proteins. The focus for determining whether applicants were in possession of claimed nucleic acids or proteins has been determined, at least in part, by considering whether the applicant has: (1) established that one skilled in the art in possession of the deposited clone would have been aware of both the

DNA sequence and the encoded amino acid sequence, or would be able to determine these sequences without undue experimentation, (2) established that the DNA and amino acid sequences are described in a manner such that one skilled in the art could distinguish them from other sequences, and (3) resequenced a clone which is identical to that of the deposit and established a "chain of custody" for this clone. See e.g., *Ex parte Maizel*, 27 U.S.P.Q.2d 1662, 1669-1670 (B.P.A.I. 1992).

Submitted herewith is a Declaration of Craig Rosen Under 37 CFR §1.132 (the "Rosen Declaration") which describes certain events involving the characterization of the nucleotide sequence of cDNA clone HE2CA82 which encodes ICE-LAP 3. The HE2CA82 cDNA clone was deposited with the American Type Culture Collection (ATCC) on August 25, 1994; i.e., prior to the filing date of parent application US Serial No. 08/334,251 (filed November 1, 1994). Exhibit A of the Rosen Declaration is a copy of the contract for ATCC Deposit No. 75875. The Examiner will note that the present specification states on page 5, lines 7 and 8, that ATCC Deposit No. 75875 contains the cDNA which encodes ICE-LAP 3.

The corrected nucleic acid and deduced amino acid sequences (shown in replacement Figures 1 and 3 submitted herewith) were determined by reanalyzing cDNA clone HE2CA82 (ATCC Accession No. 75875); i.e., the same cDNA clone from which the originally presented sequences were determined (see paragraph 4 of the Rosen Declaration). The corrected sequence information was published by the present inventors in a peer-reviewed scientific journal article (see Exhibit B of the Rosen Declaration).

The analysis needed to determine the complete and correct sequences of cDNA clone HE2CA82 were well within the skill of the ordinary artisan as of the filing date of the parent application US Serial No. 08/334,251 (filed November 1, 1994), and such analysis would not have required undue experimentation (see paragraph 5 of the Rosen Declaration).

Exhibit D of the Rosen Declaration shows the four (4) nucleotide differences between the original sequence ("PF140" in the Exhibit) and the corrected sequence ("Duan" in the Exhibit).

In summary, because Applicants have demonstrated that the corrected sequences are inherently present in the deposited material, and because Applicants have demonstrated "chain of custody" for the material originally sequenced and the resequenced material, correction of the originally presented sequence information herein is not new matter.

The specification was amended on page 5 to correct the location of the pentapeptide QACRG in SEQ ID NO:2.

The specification was further amended on page 5 to correct the spelling of "homolog".

Claims 61, 64, and 65 were amended herein. Support for the amended claims can be found in the specification and claims as originally filed. In particular, support for claims 61(b) and 64 which recite a polypeptide "lacking an N-terminal methionine residue" can be found in the specification, for example, on page 17, lines 32 and 33.


Accordingly, no new matter has been added herein.

VII. Summary

Applicants believe that the application is in condition for allowance and a notice to that effect is earnestly solicited. If the Examiner believes that a telephone conference would help to facilitate prosecution of the present application she is encouraged to call Applicants' representative at the telephone number listed below.

Respectfully submitted,

Dated: 3/22/99



A. Anders Brookes (Reg. No. 36,373)
Attorney for Applicants

Human Genome Sciences, Inc.
9410 Key West Avenue
Rockville, MD 20850
(301) 610-5761 (phone)

AAB/mbp

0985863-070204

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Application of: He et al.

Attorney Docket No.: PF140C2

Application Serial No.: Unassigned

Art Unit: Unassigned

Filed: Herewith

Examiner: Unassigned

Title: Interleukin-1 Beta Converting Enzyme
Like Apoptosis Protease-3 and 4

VERSION OF AMENDMENTS WITH MARKINGS
TO SHOW CHANGES MADE

In the Specification:

Paragraph beginning at page 4, line 7, has been amended as follows:

Figures 1A-B shows the cDNA and corresponding deduced amino acid sequence of ICE-LAP-3. The polypeptide encoded by the amino acid sequence shown is the putative mature form of the polypeptide (minus the initial methionine residue), and the standard one-letter abbreviation for amino acids is used.

Paragraph beginning at page 4, line 12, has been amended as follows:

Figures 2A-B shows the cDNA and corresponding deduced amino acid sequence of ICE-LAP-4. The polypeptide encoded by the amino acid sequence shown is the putative mature form of the polypeptide (minus the initial methionine residue).

Paragraph beginning at page 4, line 16, has been amended as follows:

Figures 3A-C shows an amino acid sequence comparison between ICE-LAP-3, ICE-LAP-4, human ICE and the C. elegans cell death gene ced-3. Shaded areas represent amino acid matches between the different sequences.

Paragraph beginning at page 4, line 20, has been amended as follows:

098553-070204

Sequencing inaccuracies are a common problem when attempting to determine polynucleotide sequences. Accordingly, the sequences of Figures 1A-B and 2A-B are based on several sequencing runs and the sequencing accuracy is considered to be at least 97%.

Paragraph beginning at page 4, line 25, has been amended as follows:

In accordance with an aspect of the present invention, there are provided isolated nucleic acids (polynucleotides) which encode the mature polypeptides having the deduced amino acid sequence of Figures 1A-B and 2A-B or for the mature polypeptide encoded by the cDNA of the clones deposited as ATCC Deposit No. 75875 encoding ICE-LAP-3, and ATCC Deposit No. 75873 encoding ICE-LAP-4, which were deposited August 25, 1994.

Paragraph beginning at page 4, line 33, has been amended as follows:

The polynucleotide encoding ICE-LAP-3 can be detected from human prostate, human endometrial tumor, human pancreatic tumor, human adrenal gland tumor and human tonsil. The full-length encoding ICE-LAP-3 was discovered in a cDNA library derived from human endometrial tumor. It is structurally related to the Interleukin-1 converting enzyme family. It contains an open reading frame encoding a protein of approximately 341 amino acid residues. The protein exhibits the highest degree of homology to *C. elegans* cell death gene *ced-3* which is a ~~homolog~~ homolog of human interleukin-1 converting enzyme, with 68 % similarity and 43 % identity over the entire amino acid sequence. It should be pointed out that the pentapeptide QACRG is conserved and is located at amino acid position ~~259-263~~ 184-188.

Paragraph beginning at page 5, line 21, has been amended as follows:

The polynucleotides of the present invention may be in the form of RNA or in the form of DNA, which DNA includes cDNA, genomic DNA, and synthetic DNA. The DNA may be double-stranded or single-stranded, and if single stranded may be the coding strand or non-coding (anti-sense) strand. The coding sequence which encode the mature polypeptides may be identical to the coding sequence shown in Figures 1A-B and 2A-B or that of the deposited clones or may be a different coding sequence which coding sequence, as a

result of the redundancy or degeneracy of the genetic code, encode the same mature polypeptides, and derivatives thereof, as the DNA of Figures 1A-B and 2A-B or the deposited cDNA.

Paragraph beginning at page 5, line 34, has been amended as follows:

The polynucleotides which encode for the mature polypeptides of Figures 1A-B and 2A-B or for the mature polypeptides encoded by the deposited cDNAs may include: only the coding sequence for the mature polypeptide; the coding sequence for the mature polypeptide and additional coding sequence; the coding sequence for the mature polypeptide (and optionally additional coding sequence) and non-coding sequence, such as introns or non-coding sequence 5' and/or 3' of the coding sequence for the mature polypeptide.

Paragraph beginning at page 6, line 10, has been amended as follows:

The present invention further relates to variants of the hereinabove described polynucleotides which encode for fragments, analogs and derivatives of the polypeptides having the deduced amino acid sequence of Figures 1A-B and 2A-B or the polypeptides encoded by the cDNA of the deposited clones. The variants of the polynucleotides may be naturally occurring allelic variants of the polynucleotides or non-naturally occurring variants of the polynucleotides.

Paragraph beginning at page 6, line 18, has been amended as follows:

Thus, the present invention includes polynucleotides encoding the same mature polypeptides as shown in Figures 1A-B and 2A-B or the same mature polypeptides encoded by the cDNA of the deposited clones as well as variants of such polynucleotides which variants encode for a fragment, derivative or analog of the polypeptides of Figures 1A-B and 2A-B or the polypeptides encoded by the cDNA of the deposited clones. Such nucleotide variants include deletion variants, substitution variants and addition or insertion variants.

Paragraph beginning at page 6, line 27, has been amended as follows:

As hereinabove indicated, the polynucleotides may have a coding sequence which is a naturally occurring allelic variant of the coding sequence shown in

Figures 1A-B and 2A-B or of the coding sequence of the deposited clones. As known in the art, an allelic variant is an alternate form of a polynucleotide sequence which may have a substitution, deletion or addition of nucleotides, which does not substantially alter the function of the encoded polypeptides. The polynucleotides may also encode for a proprotein which is the mature protein plus additional 5' amino acid residues.

Paragraph beginning at page 7, line 11, has been amended as follows:

The present invention further relates to polynucleotides which hybridize to the hereinabove-described sequences if there is at least 70%, preferably at least 90%, and more preferably at least 95% identity between the sequences. The present invention particularly relates to polynucleotides which hybridize under stringent conditions to the hereinabove-described polynucleotides. As herein used, the term "stringent conditions" means hybridization will occur only if there is at least 95% and preferably at least 97% identity between the sequences. The polynucleotides which hybridize to the hereinabove described polynucleotides in a preferred embodiment encode polypeptides which either retain substantially the same biological function or activity as the mature polypeptide encoded by the cDNAs of Figures 1A-B and 2A-B or the deposited cDNA(s).

Paragraph beginning at page 8, line 4, has been amended as follows:

The present invention further relates to ICE-LAP-3 and 4 polypeptides which have the deduced amino acid sequence of Figures 1A-B and 2A-B or which has the amino acid sequence encoded by the deposited cDNAs, as well as fragments, analogs and derivatives of such polypeptides.

Paragraph beginning at page 8, line 9, has been amended as follows:

The terms "fragment," "derivative" and "analog" when referring to the polypeptides of Figures 1A-B and 2A-B or that encoded by the deposited cDNA, means polypeptides which retain essentially the same biological function or activity as such polypeptides, and wherein derivatives include

polypeptides with enhanced or reduced biological function. An analog includes a proprotein portion to produce active mature polypeptides.

Paragraph beginning at page 8, line 20, has been amended as follows:

The fragment, derivative or analog of the polypeptides of Figures 1A-B and 2A-C or that encoded by the deposited cDNAs may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one in which the additional amino acids are fused to the mature polypeptide for purification of the mature polypeptide. Such fragments, derivatives and analogs are deemed to be within the scope of those skilled in the art from the teachings herein.

Paragraph beginning at page 26, line 11, has been amended as follows:

--The DNA sequence encoding ICE-LAP-3, ATCC # 75875, is initially amplified using PCR oligonucleotide primers corresponding to the 5' sequences of the processed ICE-LAP-3 protein (minus the signal peptide sequence) and the vector sequences 3' to the ICE-LAP-3 gene. Additional nucleotides corresponding to ICE-LAP-3 are added to the 5' and 3' sequences respectively. The 5' oligonucleotide primer has the sequence 5' GATCGGATCCATGCGTGCGGGGACACGGGTC 3' (SEQ ID NO:5) contains a Bam HI restriction enzyme site (underlined) followed by 18 nucleotides of ICE-LAP-3 coding sequence starting from the presumed terminal amino acid of the processed protein codon. The 3' sequence 5' GTACTCTAGATCATTACCCCTGGTGGAGGAT 3' (SEQ ID NO:6) contains complementary sequences to an Xba I site (underlined) followed by 21 nucleotides of ICE-LAP-3. The restriction enzyme sites correspond to the restriction enzyme sites on the bacterial expression vector pQE-9 (Qiagen, Inc. 9259 Eton Avenue, Chatsworth, CA, 91311). pQE-9 encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable

0985263 070201
T02020 0935360

promoter operator (P/O), a ribosome binding site (RBS), a 6-His tag and restriction enzyme sites. PQE-9 is then digested with Bam HI and Xba I. The amplified sequences are ligated into pQE-9 and are inserted in frame with the sequence encoding for the histidine tag and the RBS. The ligation mixture is then used to transform *E. coli* available from Qiagen under the trademark M15/rep 4 by the procedure described in Sambrook, J. et al., Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989). M15/rep4 contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis. Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG ("Isopropyl-B-D-thiogalactopyranoside") is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression. Cells are grown an extra 3 to 4 hours. Cells are then harvested by centrifugation. The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl. After clarification, solubilized ICE-LAP-3 is purified from this solution by chromatography on a Nickel-Chelate column under conditions that allow for tight binding by proteins containing the 6-His tag (Hochuli, E. et al., J. Chromatography 411:177-184 (1984)). ICE-LAP-3 (95% pure) is eluted from the column in 6 molar guanidine HCl pH 5.0 and for the purpose of renaturation adjusted to 3 molar guanidine HCl, 100mM sodium phosphate, 10 mmolar glutathione (reduced) and 2 mmolar glutathione (oxidized). After incubation in this solution for 12 hours the protein was dialyzed to 10 mmolar sodium phosphate.—

Paragraph beginning at page 28, line 1, has been amended as follows:

-- The DNA sequence encoding ICE-LAP-4, ATCC # 75873, is initially amplified using PCR oligonucleotide primers corresponding to the 5' sequences of the processed ICE-LAP-4 protein (minus the signal peptide

sequence) and the vector sequences 3' to the ICE-LAP-4 gene. Additional nucleotides corresponding to ICE-LAP-4 are added to the 5' and 3' sequences respectively. The 5' oligonucleotide primer has the sequence 5' GATCGGATCCATGGAGAACTGAAACTCA 3' (SEQ ID NO:7) contains a Bam HI restriction enzyme site (underlined) followed by 18 nucleotides of ICE-LAP-3 coding sequence starting from the presumed terminal amino acid of the processed protein codon. The 3' sequence 5' GTACTCTAGATTAGTGATAAAAATAGAGTTC 3' (SEQ ID NO:8) contains complementary sequences to an Xba I site (underlined) followed by 21 nucleotides of ICE-LAP-4. The restriction enzyme sites correspond to the restriction enzyme sites on the bacterial expression vector pQE-9 (Qiagen, Inc. 9259 Eton Avenue, Chatsworth, CA, 91311). pQE-9 encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter operator (P/O), a ribosome binding site (RBS), a 6-His tag and restriction enzyme sites. pQE-9 is then digested with Bam HI and Xba I. The amplified sequences are ligated into pQE-9 and are inserted in frame with the sequence encoding for the histidine tag and the RBS. The ligation mixture is then used to transform *E. coli* available from Qiagen under the trademark M15/rep 4 by the procedure described in Sambrook, J. et al., Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989). M15/rep4 contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis. Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG ("Isopropyl-B-D-thiogalactopyranoside") is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression. Cells are grown an extra 3 to 4 hours. Cells are then harvested by centrifugation. The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl. After clarification, solubilized ICE-LAP-4 is purified from

102040 025650

this solution by chromatography on a Nickel-Chelate column under conditions that allow for tight binding by proteins containing the 6-His tag (Hochuli, E. et al., J. Chromatography 411:177-184 (1984)). ICE-LAP-4 (95% pure is eluted from the column in 6 molar guanidine HCL pH 5.0 and for the purpose of renaturation adjusted to 3 molar guanidine HCl, 100mM sodium phosphate, 10 mmolar gluathione (reduced) and 2 mmolar glutathione (oxidized). After incubation in this solution for 12 hours the protein was dialyzed to 10 mmolar sodium phosphate.--

Paragraph beginning at page 30, line 7, has been amended as follows:

-- The DNA sequence encoding for ICE-LAP-3, ATCC # 75875, was constructed by PCR on the full-length ICE-LAP-3 using two primers: the 5' primer 5' GACTATGCGTGC GGGGACACGG 3' (SEQ ID NO:9) contains the ICE-LAP-3 translational initiation site ATG followed by 5 nucleotides of ICE-LAP-3 coding sequence starting from the initiation codon; the 3' sequence 5' AATCAAGCGTAGTCTGGGACGTCGTATGGGTATTCACCCTGGTGGA GGATTG 3' (SEQ ID NO:10) contains translation stop codon, HA tag and the last 21 nucleotides of the ICE-LAP-3 coding sequence (not including the stop codon). Therefore, the PCR product contains the ICE-LAP-3 coding sequence followed by HA tag fused in frame, and a translation termination stop codon next to the HA tag. The PCR amplified DNA fragment was ligated with pcDNA1/Amp by blunt end ligation. The ligation mixture was transformed into E. coli strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037) the transformed culture was plated on ampicillin media plates and resistant colonies were selected. Plasmid DNA was isolated from transformants and examined by restriction analysis for the presence of the correct fragment. For expression of the recombinant ICE-LAP-3, COS cells were transfected with the expression vector by DEAE-DEXTRAN method (J. Sambrook, E. Fritsch, T. Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989)). The expression of the ICE-LAP-3 HA protein was detected by radiolabelling and immunoprecipitation method (E. Harlow, D. Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory

Press, (1988)). Cells were labelled for 8 hours with ³⁵S-cysteine two days post tranfection. Culture media were then collected and cells were lysed with detergent (RIPA buffer (150 mM NaCl, 1% NP-40, 0.1% SDS, 1% NP-40, 0.5% DOC, 50mM Tris, pH 7.5) (Wilson, I. Et al., Id. 37:767 (1984)). Both cell lysate and culture media were precipitated with a HA specific monoclonal antibody. Proteins precipitated were analyzed on 15% SDS-PAGE gels.--

Paragraph beginning at page 32, line 20, has been amended as follows:

-- The DNA sequence encoding for ICE-LAP-4, ATCC # 75873, was constructed by PCR on the full-length ICE-LAP-4 using two primers: the 5' primer 5' ACCATGGAGAACACTGAAAAC 3' (SEQ ID NO:11) contains the ICE-LAP-4 translational initiation site ATG followed by 5 nucleotides of ICE-LAP-4 coding sequence starting from the initiation codon; the 3' sequence 5'

AATCAAGCGTAGTCTGGGACGTCGGTATGGGTAGTGATAAAAATA GAGTTCTTT 3' (SEQ ID NO:12) contains translation stop codon, HA tag and the last 21 nucleotides of the ICE-LAP-4 coding sequence (not including the stop codon). Therefore, the PCR product contains the ICE-LAP-4 coding sequence followed by HA tag fused in frame, and a translation termination stop codon next to the HA tag. The PCR amplified DNA fragment was ligated with pcDNAI/Amp by blunt end ligation. The ligation mixture was transformed into E. coli strain SURE (available from Stratagene Cloning Systems, 11099 North Torrey Pines Road, La Jolla, CA 92037) the transformed culture was plated on ampicillin media plates and resistant colonies were selected. Plasmid DNA was isolated from transformants and examined by restriction analysis for the presence of the correct fragment. For expression of the recombinant ICE-LAP-4, COS cells were transfected with the expression vector by DEAE-DEXTRAN method (J. Sambrook, E. Fritsch, T. Maniatis, Molecular Cloning: A Laboratory Manual, Cold Spring Laboratory Press, (1989)). The expression of the ICE-LAP-4 HA protein was detected by radiolabelling and immunoprecipitation method (E. Harlow, D. Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, (1988)). Cells were labelled for 8 hours with ³⁵S-cysteine two days post tranfection. Culture media were then collected and cells were lysed with

detergent (RIPA buffer (150 mM NaCl, 1% NP-40, 0.1% SDS, 1% NP-40, 0.5% DOC, 50mM Tris, pH 7.5) (Wilson, I. Et al., Id. 37:767 (1984))). Both cell lysate and culture media were precipitated with a HA specific monoclonal antibody. Proteins precipitated were analyzed on 15% SDS-PAGE gels.--

09259360

1 GCACGAGAACTTTGCTGTGCGCGTTCTCCCGCGCGCGGGCTCAACTTTGTAGAGCGAGG

61 GGCCAACCTTGGCAGAGCGCGCGGCCAGCTTTGCAGAGAGCGCCCTCCAGGGACTATGCGT

1 1 ~~M R~~

121 GCGGGGACACGGGTCGCTTTGGGCTCTTCCACCCCTGCGGAGCGCACTACCCCGAGCCAG

3 ~~A C T R V A L G S S T P A E R T T P S Q~~

181 GGGCGGTGCAAGCCCCGCGGCCCTACCCAGGGCGGCTCCTCCCTCCGCAGCGCCGAGA

23 ~~G R C K P R P A L P R A A P P S A A P R~~

241 CTTTTAGTTTCGCTTTTCGCTAAAGGGGCCCCAGACCCTTGCTGCGGAGCGACGGAGAGAG

43 ~~L L V S L S L K G P Q T L A A E R R R E~~

301 ACTGTGCCAGTCCCAGCCGCCCTACCGCCGTGGGAACGATGGCAGATGATCAGGGCTGT

63 ~~T V P V P A A L P P W E R~~ ~~M~~ ~~Q~~ ~~A~~ ~~N~~ ~~D~~ ~~I~~ ~~D~~ ~~Q~~ ~~G~~ ~~C~~

361 ATTGAAGAGCAGGGGGTTGAGGATTCAGCAAATGAAGATTGATGGATGCTAAGCCGAC

83 I E E Q G V E D S A N E D S V D A K P D

421 CGGTCCTCGTTTGTACCGTCCCTCTTCAGTAAGAAGAAGAAAAATGTCACCATGCGATCC

103 R S S S F V P S L F S K K K K N V T M R S

481 ATCAAGACCACCCGGGACCGAGTGCCTACATATCAGTACAACATGAATTTTGAAGCTG

123 I K T T R D R V P T Y Q Y N M N F E K L

541 GGCAAATGCATCATAATAAACAACAAGAACTTTGATAAGTGACAGGTATGGGCGTTTCCA

143 G K C I I I N N K N F D K V T G M G V R

601 AACGGAACAGACAAAGATGCCGAGGCGCTCTTCAAGTGCTTCCGAAGCCTGGGTTTGTAC

163 N G T D K D A E A L F K C F R S L G F D

661 GTGATTGTCTATAATGACTGCTCTTGTGCCAAGATGCAAGATCTGCTTAAAAAGCTTCT

183 V I V Y N D C S C A K M Q D L L K K A S

721 GAAGAGGACCATACAAATGCCGCTGCTTCGCCTGCATCCTCTTAAGCCATGGAGAAGAA

203 E E D H T N A A C F A C I L L S H G E E

781 AATGTAATTTATGGAAAGATGGTGTACACCAATAAAGGATTTGACAGCCCACTTTAGG

223 N V I Y G K D G V T C P I K D L T A H F R

841 GGGGATAGATGCAAAACCCTTTTAGAGAAACCCAAACTCTTCTTCATTACGGCTTGCCGA

243 G D R C K T L L E K P K L F F I Q A C R

901 GGGACCGAGCTTGATGATGATCCATCCAGGCCGACTCGGGGCCCATCAATGACACAGATGCT

263 G T E L D D ~~A~~ I Q A D S G P I N D T D A

961 AATCCTCGATACAAGATCCCAGTGGAAGCTGACTTCCTCTTCGCCTATTCCACGGTTCCA

283 N P R Y K I P V E A D F L F A Y S T V P

1021 GGCTATTACTCGTGAGGAGCCCAGGAAGAGGCTCCTGGTTTGTGCAAGCCCTCTGCTCC

303 G Y Y S W R S P G R G S W F V Q A L C S

1081 ATCCTGGAGGAGCACGGAAAAGACCTGGAATCATGCAATCCTCACCAGGGTGAATGA

323 I L E E H G K D L E I M Q I L ~~H~~ ~~T~~ ~~R~~ ~~V~~ ~~N~~ ~~D~~

1141 CAGAGTTGCCAGGCACCTTTGAGTCTCAGTCTGATGACCCACACTTCCATGAGAAGAAGCA

~~R V A R H F E S Q S D D P H F H E K K Q~~

1201 GATCCCTGTGTGGTCTCCATGCTCACCAAGGAAGCTTACTTCACTCAATAGCCATATCA

~~T P C V V S N L T K E L Y F S Q~~

1261 GGGGTACATTCTAGCTGAGAAGCAATGGGTCACTCATTAAATGAATCACATTTTTTTATGC

1321 TCTTGAAATATTTCAGAAATTCTCCAGGATTTTAATTTTCAGGAAAATGTATT

Figure 1

1 GCACGAGCGGATGGGTGCTATTGTGAGGCGGTTGTAGAAGAGTTTCGTGAGTGCTCGCAG
 61 CTCATACCTGTGGCTGTGTATCCGTGGCCACAGCTGGTTGGCGTCGCCTTGAAATCCCAG
 121 GCCGTGAGGAGTTAGCGAGCCCTGCTCACACTCGGCGCTCTGGTTTTTCGGTGGGTGTGCC
 181 CTGCACCTGCCTCTTCCCGCATTCTCATTAAATAAAGGTATCCATGGAGAACACTGAAAAC
 1 M E N T E N
 241 TCAGTGGATTCAAAATCCATTAATAAATTTGGAACCAAAGATCATAACATGGAAGCGAATCA
 7 S V D S K S I K N L E P K I I H G S E S
 301 ATGGACTCTGGAATATCCCTGGACAACAGTTATAAAATGGATTATCCTGAGATGGGTTTA
 27 M D S G I S L D N S Y K M D Y P E M G L
 361 TGTATAATAATTAATAATAAGAATTTTCATAAAAGCACTGGAATGACATCTCGGTCTGGT
 47 C I I I N N K N F H K S T G M T S R S G
 421 ACAGATGTCGATGCAGCAAACCTCAGGAAACATTTCAGAAACTTGAAATATGAAGTCAGG
 67 T D V D A A N L R E T F R N L K Y E V R
 481 AATAAAATGATCTTACACGTGAAGAAATTGTGGAATTGATGCGTGATGTTTCTAAAGAA
 87 N K N D L T R E E I V E L M R D V S K E
 541 GATCACAGCAAAGGAGCAGTTTTGTTTGTGTGCTTCTGAGCCATGGTGAAGAAGGAATA
 107 D H S K R S S F V C V L L S H G E E G I
 601 ATTTTTGGAACAAATGGACCTGTTGACCTGAAAAAATAACAACTTTTTTCAGAGGGGAT
 127 I F G T N G P V D L K K I T N F F R G D
 661 CGTTGTAGAAGTCTAACTGGAACCCAACTTTTCATTATTCAGGCCTGCCGTGGTACA
 147 R C R S L T G K P K L F I I Q A C R G T
 721 GAACTGGACTGTGGCATTGAGACAGACAGTGGTGTGATGATGACATGGCGTGTCTATAA
 167 E L D C G I E T D S G V D D D M A C H K
 781 ATACCAGTGGAGGCCGACTTCTTGTATGCATACTCCACAGCACCTGGTTATTATTCTTGG
 187 I P V E A D F L Y A Y S T A P G Y Y S W
 841 CGAAATTCAAAGGATGGCTCCTGGTTCATCCAGTCGCTTTGTGCCATGCTGAAACAGTAT
 207 R N S K D G S W F I Q S L C A M L K Q Y
 901 GCCGACAAGCTTGAATTTATGCACATTCTTACCCGGGTTAACCGAAAGGTGGCAACAGAA
 227 A D K L E F M H I L T R V N R K V A T E
 961 TTTGAGTCCTTTTCCTTTGACGCTACTTTTCATGCAAAGAAACAGATTCATGTATTGTT
 247 F E S F S F D A T F H A K K Q I P C I V
 1021 TCCATGCTCACAAAAGAACTCTATTTTTATCACTAAAGAAATGGTTGGTTGGTGGTTTTT
 267 S M L T K E L Y F Y H *
 1081 TTTAGTTTGTATGCCAAGTGAGAAGATGGTATATTTGGGTACTGTATTTCCCTCTCATTG
 1141 GGGACCTACTCTCATGCTG

Figure 2

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
84